

THE CANADIAN ENTOMOLOGIST

Index to Volume 119 (1987)

SUPPLEMENTS AND MEMOIRS



THE ENTOMOLOGICAL SOCIETY OF CANADA, OTTAWA

- Akar, H. Jr., and E.A. Osgood. The method by which larvae of *Dasineura balsamicola* (Lintner) (Diptera: Cecidomyiidae) gain access to the interior of galls induced by *Paradiplosis tumifex* Gagné (Diptera: Cecidomyiidae), 395.
- Alfaro, R.I. See Otvos *et al.*, 707.
- Allen, O.B. See McAuslane *et al.*, 577.
- Allen, R.K., and C.M. Murvosh. New Baetidae from the southwestern United States and northern Mexico (Ephemeroptera: Insecta), with notes, 1095.
- Allen, W.R. See Broadbent *et al.*, 501.
- Anderson, J.R., and J.A. Shemanchuk. The biology of *Simulium arcticum* Malloch in Alberta. Part I. Obligate anaotogeny in *S. arcticum* and other black flies (Diptera: Simuliidae), 21.
- The biology of *Simulium arcticum* Malloch in Alberta. Part II. Seasonal parity structure and mermithid parasitism of populations attacking cattle and flying over the Athabasca River, 29.
- Anderson, T.E. See Hajek *et al.*, 959.
- Arthur, A. See Sharkey *et al.*, 611.
- Ankevoeld, L.S. The identity of *Donacia cuprea* Kirby, 1837, and *Donacia quadricollis* Say, 1827, with a taxonomic revision of members of the *Donacia subtilis* Kunze-group (Coleoptera: Chrysomelidae: Donaciinae), 629.
- Axtell, R.C. See Lysyk and Axtell, 427.
- Ayre, G.L. See Struble *et al.*, 291.
- See Struble *et al.*, 301.
- Bakke, A. See McLean *et al.*, 273.
- Barbosa, P., and P. Martinat. Causes and ecological implications of egg retention in the gypsy moth, *Lymantria dispar* (L.) (Lepidoptera: Lymantriidae), 765.
- Barker, P.S. See Smith and Barker, 873.
- Barker, P.S., and L.B. Smith. Spatial distribution of insect species in granary residues in the Prairie Provinces, 1125.
- Barron, J. See Sharkey *et al.*, 611.
- Baumgaertner, J.U. See Gutierrez *et al.*, 109.
- Behan-Pelletier, V.M. Redefinition of *Ametoproctus* (Acari: Oribatida) with descriptions of new species, 505.
- Beshear, R.J. See Weiss and Beshear, 399.
- Biever, K.D. See Hajek *et al.*, 959.
- Bilodeau, R.J. See Turnock *et al.*, 419.
- Bisdee, G. See Sharkey *et al.*, 611.
- Bisdee, H.E. See Wylie and Bisdee, 857.
- Boiteau, G. A red color morph of the Colorado potato beetle, *Leptinotarsa decemlineata* (Say) (Coleoptera: Chrysomelidae), 957.
- Boiteau, G., R.H. Parry, and G.R. Harris. Insecticide resistance in New Brunswick populations of the Colorado potato beetle (Coleoptera: Chrysomelidae), 459.
- Borden, J.H. See Javer *et al.*, 605.
- See Osborn and Borden, 1109.
- Bousquet, Y. Description of the larva of *Helluomorphoides praeustus bicolor* Harris with comments on the relationships of the Helluonini (Coleoptera: Carabidae), 921.
- Bracken, G.K. Relation between pod damage caused by larvae of bertha armyworm, *Mamestra configurata* Walker (Lepidoptera: Noctuidae), and yield loss, shelling, and seed quality in canola, 365.
- Broadbent, A.B., W.R. Allen, and R.G. Roottit. The association of *Frankliniella occidentalis* (Pergande) (Thysanoptera: Thripidae) with greenhouse crops and the tomato spotted wilt virus in Ontario, 501.
- Buckle, D.J. See Leech and Buckle, 1145.
- Burgess, L., and C.F. Hinks. Predation on adults of the crucifer flea beetle, *Phyllotreta cruciferae* (Goeze), by the northern fall field cricket, *Gryllus pennsylvanicus* Burmeister (Orthoptera: Gryllidae), 495.
- Byers, J.R. See Struble *et al.*, 291.
- See Struble *et al.*, 301.
- Byers, J.R., and D.L. Struble. Monitoring population levels of eight species of noctuids with sex-attractant traps in southern Alberta, 1978-1983: specificity of attractants and effect of target species abundance, 541.
- Byers, J.R., D.L. Struble, and G.B. Schaalje. Monitoring populations levels of eight species of noctuids with sex-attractant traps in southern Alberta, 1978-1983: consistency of trap catches and precision of population estimates, 557.
- Campbell, J.M. *Anthobioides pubescens*, an unusual new genus and species of Omaliinae (Coleoptera: Staphylinidae) from Washington, 1027.
- Campbell, J.M., and D.S. Chandler. *Omalorphanus aenigma*, an unusual new genus and species of Omaliinae (Coleoptera: Staphylinidae) from Oregon, 315.
- Canaday, C.L. Comparison of insect fauna captured in six different trap types in a Douglas-fir forest, 1101.
- Cannings, S.G. *Capnia licina* Jewett (Plecoptera: Capniidae): a description of the adult female and redescription of the male, 439.
- Capinera, J.L. See Horton and Capinera, 729.

- Capinera, J.L., and D.C. Thompson. Dynamics and structure of grasshopper assemblages in shortgrass prairie, 567.
- Carrow, J.R. See Smith *et al.*, 251.
- Chandler, D.S. See Campbell and Chandler, 315.
- Charlet, L.D. Seasonal dynamics of the sunflower stem weevil, *Cylindrocaptus adpersus* (LeConte) (Coleoptera: Curculionidae), on cultivated sunflower in the northern great plains, 1133.
- Chianese, R. See Lashomb *et al.*, 75.
- Cloutier, C. See Morris and Cloutier, 381.
- Coderre, D., L. Provencher, and J.-C. Tourneur. Oviposition and niche partitioning in aphidophagous insects on maize, 195.
- Cohen, M.B., and M. Mackauer. Intrinsic rate of increase and temperature coefficients of the aphid parasite *Ephedrus californicus* Baker (Hymenoptera: Aphididae), 231.
- Corkum, L.D. Patterns in mayfly (Ephemeroptera) wing length: adaptation to dispersal?, 783.
- Cornet, M. See Young *et al.*, 481.
- Cunningham, J.C. See Orvos *et al.*, 697.
- See Orvos *et al.*, 707.
- Dawson, A.F. See Hulme *et al.*, 751.
- Dondale, C.D., and J.H. Redner. The *atrata*, *cubana*, *ferruginea*, *moesta*, *monticola*, *saltuaria*, and *solitaria* groups of the spider genus *Pardosa* in North America (Araneae: Lycosidae), 1.
- Dondale, L.M., and D.M. Lehmkuhl. Stoneflies (Plecoptera) of the Lake Athabasca region of northern Saskatchewan and their biogeographical affinities, 1059.
- Eidt, D.C. See Fairchild *et al.*, 647.
- Ellis, C.K. See Gutierrez *et al.*, 109.
- Ellis, C.R. See McAuslane *et al.*, 577.
- Erman, N.A. See Wiggins and Erman, 867.
- Ewen, A.B. See Germida *et al.*, 355.
- Ewen, A.B., and M.K. Mukerji. Field evaluation of carbofuran bait against grasshopper (Orthoptera: Acrididae) populations in Saskatchewan, 537.
- Fairchild, W.L., D.C. Eidt, and C.A.A. Weaver. Effects of fenitrothion insecticide on inhabitants of leaves of the pitcher plant, *Sarracenia purpurea* L., 647.
- Fairchild, W.L., and D.J. Lewis. Parasitic water mite larvae (Hydrachnida: Hygrobatidae) associated with caddisfly larvae (Trichoptera: Leptoceridae, Limnephilidae), 809.
- Ferran, A. See Lapchin *et al.*, 815.
- Ferro, D.N. See Hajek *et al.*, 959.
- Finnamore, A.T. A new genus and species of psenine wasp from Africa and a key to genera (Hymenoptera: Pemphredonidae: Pseninae), 1081.
- Footitt, R.G. See Broadbent *et al.*, 501.
- Friskie, L.M. See Orvos *et al.*, 697.
- Gerber, G.H. Reproductive cycles of the red turnip beetle, *Entomoscelis americana* Brown (Coleoptera: Chrysomelidae), 1069.
- Germida, J.J., A.B. Ewen, and E.E. Onofriechuk. *Nosema locustae* Canning (Microsporidia) spore populations in treated field soils and resident grasshopper populations, 355.
- Gilleson, L.A. A note on fecundity of the aphid predator, *Aphidoletes aphidimyza* (Rondani) (Diptera: Cecidomyiidae), 1147.
- Goettel, M.S. Serial *in vivo* passage of the entomopathogenic hyphomycete *Tolypocladium cylindrosporum* in mosquitoes, 599.
- Gordon, R. See Young *et al.*, 481.
- Graham, K. See Syed and Graham, 489.
- Guppy, J.C., and F. Meloche. Life history and description of the immature stages of *Dacnusa dryas* (Nixon) (Hymenoptera: Braconidae), a European parasite of the alfalfa blotch leafminer (Diptera: Agromyzidae) in eastern Canada, 281.
- Gutierrez, A.P., F. Schulthess, L.T. Wilson, A.M. Villacorta, C.K. Ellis, and J.U. Baumgaertner. Energy acquisition and allocation in plants and insects: a hypothesis for the possible role of hormones in insect feeding patterns, 109.
- Hagley, E.A.C., and N. Miles. Release of *Chrysoperla carnea* Stephens (Neuroptera: Chrysopidae) for control of *Tetranychus urticae* Koch (Acarina: Tetranychidae) on peach grown in a protected environment structure, 205.
- Hajek, A.E., R.S. Soper, D.W. Roberts, T.E. Anderson, K.D. Biever, D.N. Ferro, R.A. LeBrun, and R.H. Storch. Foliar applications of *Beauveria bassiana* (Balsamo) Vuillemin for control of the Colorado potato beetle, *Leptinotarsa decemlineata* (Say) (Coleoptera: Chrysomelidae): an overview of pilot test results from the northern United States, 959.
- Hamilton, K.G.A., and D.W. Langor. Leafhopper fauna of Newfoundland and Cape Breton islands (Rhynchota: Homoptera: Cicadellidae), 663.

- Harris, G.R. See Boiteau *et al.*, 459.
Harris, J.W.E. See Hulme *et al.*, 751.
Hidaka, T. See Obata and Hidaka, 603.
Hinks, C.F. See Burgess and Hinks, 495.
Hogg, D.B. See Thiboldeaux *et al.*, 1055.
Horton, D.R., and J.L. Capinera. Seasonal and host plant effects on parasitism of Colorado potato beetle by *Myiophorus doryphorae* (Riley) (Diptera: Tachinidae), 729.
Hubbes, M. See Smith *et al.*, 251.
Huber, J.T. Review of *Schizophragma* Ogloblin and the non-Australian species of *Stethynium* Enoch (Hymenoptera: Mymaridae), 823.
Hulme, M.A., J.W.E. Harris, and A.F. Dawson. Exploiting adult girth to separate *Pissodes strobi* (Peck) (Coleoptera: Curculionidae) from associated insects in leaders of *Picea sitchensis* (Bong.) Carr., 751.
Hutchison, W.D. See Thiboldeaux *et al.*, 1055.
Iperti, G. See Lapchin *et al.*, 815.
Jansson, R.K. See Lashomb *et al.*, 75.
Jaques, R.P., J.E. Laing, D.R. Laing, and D.S.K. Yu. Effectiveness and persistence of the granulosis virus of the codling moth *Cydia pomonella* (L.) (Lepidoptera: Olethreutidae) on apple, 1063.
Javer, A., A.D. Wynne, J.H. Borden, and G.J.R. Judd. Pine oil: an oviposition deterrent for the onion maggot, *Delia antiqua* (Meigen) (Diptera: Anthomyiidae), 605.
Johnson, J.W. A revision of the species of *Praon* Haliday in North America north of Mexico (Hymenoptera: Aphididae), 999.
Judd, G.J.R. See Javer *et al.*, 605.
Kemp, W.P., and N.E. Sánchez. Differences in post-diapause thermal requirements for eggs of two rangeland grasshoppers, 653.
Kettela, E.G. See Turgeon *et al.*, 361.
Khachatourians, G.G. See Marcandier and Khachatourians, 893.
 See Marcandier and Khachatourians, 901.
Kononenko, V.S. See Lafontaine *et al.*, 931.
Koponen, S., and M.S. Wasbauer. A note on the biology of *Anoplus tenuicornis* (Tournier) in North America (Hymenoptera: Pompilidae), 103.
Krainacker, D. See Lashomb *et al.*, 75.
Krzeminski, W., and H.J. Teskey. New taxa of Limoniidae (Diptera: Nematocera) from Canadian amber, 887.
Kveder, T.M. See Whitfield *et al.*, 859.
Lafontaine, J.D., K. Mikkola, and V.S. Kononenko. *Anarta cordigera* (Thunberg) (Lepidoptera: Noctuidae: Hadeninae), a species complex, 931.
Laing, D.R. See Jaques *et al.*, 1063.
Laing, J.E. See Jaques *et al.*, 1063.
Lamb, R.J. See Turnock *et al.*, 419.
Langor, D.W. See Hamilton and Langor, 663.
Langor, D.W., and A.G. Raske. Emergence, host attack, and overwintering behavior of the eastern larch beetle, *Dendroctonus simplex* LeConte (Coleoptera: Scolytidae), in Newfoundland, 975.
 Reproduction and development of the eastern larch beetle, *Dendroctonus simplex* LeConte (Coleoptera: Scolytidae), in Newfoundland, 985.
Lanier, G.N. The validity of *Ips cribricollis* (Eich.) (Coleoptera: Scolytidae) as distinct from *I. grandicollis* (Eich.) and the occurrence of both species in Central America, 179.
 See Phillips *et al.*, 465.
Lapchin, L., A. Ferran, G. Iperti, J.M. Rabasse, and J.P. Lyon. Coccinellids (Coleoptera: Coccinellidae) and syrphids (Diptera: Syrphidae) as predators of aphids in cereal crops: a comparison of sampling methods, 815.
Lashomb, J., D. Krainacker, R.K. Jansson, J.S. Ng, and B. Chianese. Parasitism of *Leptinotarsa decemlineata* (Say) eggs by *Edovum puttleri* Grissell (Hymenoptera: Eulophidae): effects of host age, parasitoid age, and temperature, 75.
LeBrun, R.A. See Hajek *et al.*, 959.
Legner, E.F. Inheritance of gregarious and solitary oviposition in *Muscidifurax raptorellus* Kogan and Legner (Hymenoptera: Pteromalidae), 791.
Lee, D.A., and J.R. Spence. Developmental adaptation of the European corn borer (*Ostrinia nubilalis* Hübner) in Alberta, 371.
Leech, R., and D.J. Buckle. The first records of *Dolomedes striatus* Giebel in Alberta and Saskatchewan (Araneida: Pisauridae), 1145.
Legner, E.F. Transfer of thelytoky to arrhenotokous *Muscidifurax raptor* Girault and Sanders (Hymenoptera: Pteromalidae), 265.
Lehmkuhl, D.M. See Whiting and Lehmkuhl, 405.
 See Whiting and Lehmkuhl, 409.
 See Dossdall and Lehmkuhl, 1059.

- Lewis, D.J. See Fairchild and Lewis, 809.
- Lim, K.P. See West *et al.*, 449.
- Lindquist, E.E., and K.W. Wu. First record of the mite family Podacarinae (Acari: Mesostigmata) in Canada, with notes on other records in North America, 779.
- Lochiavo, S.R., and N.D.G. White. The loss of a grain fumigant mixture of carbon tetrachloride and carbon disulfide through sealing and various covering material, 595.
- Lyon, J.P. See Lapchin *et al.*, 815.
- Lysyk, T.J., and R.C. Axtell. A simulation model of house fly (Diptera: Muscidae) development in poultry manure, 427.
- Mackauer, M. See Cohen and Mackauer, 231.
- Maier, C.T., and D.W. Webb. Spatial and temporal variation in abundance, sex ratio, and species composition of Connecticut Rhagionidae (Diptera), 55.
- Marcandier, S., and G.G. Khachatourians. Evolution of relative humidity and temperature within a closed chamber used for entomological studies, 893.
- Susceptibility of the migratory grasshopper, *Melanoplus sanguinipes* (Fab.) (Orthoptera: Acrididae), to *Beauveria bassiana* (Bals.) Vuillemin (Hyphomycete): influence of relative humidity, 901.
- Martinat, P. See Barbosa and Martinat, 765.
- McAuslane, H.J., C.R. Ellis, and O.B. Allen. Sequential sampling of adult northern and western corn rootworms (Coleoptera: Chrysomelidae) in southern Ontario, 577.
- McLean, J.A. See Shore *et al.*, 131.
- See Sweeney and McLean, 587.
- McLean, J.A., A. Bakke, and H. Niemeyer. An evaluation of three traps and two lures for the ambrosia beetle *Trypodendron lineatum* (Oliv.) (Coleoptera: Scolytidae) in Canada, Norway, and West Germany, 273.
- McLeod, D.G.R. See Struble *et al.*, 291.
- Meighen, E.A. See Sanders and Meighen, 305.
- Meloche, F. See Guppy and Meloche, 281.
- Meyerdirk, D.E., and M.S. Moratorio. *Circulifer tenellus* (Baker), the beet leafhopper (Homoptera: Cicadellidae): laboratory studies on fecundity and longevity, 443.
- Morphometric study of *Anagrus giraulti* Crawford (Hymenoptera: Mymaridae) reared from eggs of three different cicadellid hosts, 909.
- Mikkola, K. See Lafontaine *et al.*, 931.
- Miles, N. See Hagley and Miles, 205.
- Moratorio, M.S. See Meyerdirk and Moratorio, 443.
- See Meyerdirk and Moratorio, 909.
- Morris, D.E., and C. Cloutier. Biology of the predatory fly *Coenosia tigrina* (Fab.) (Diptera: Anthomyiidae): reproduction, development, and larval feeding on earthworms in the laboratory, 361.
- Morris, O.N. Evaluation of the nematode, *Steinernema feltiae* Filipjev, for the control of the crucifer flea beetle, *Phyllotreta cruciferae* (Goeze) (Coleoptera: Chrysomelidae), 95.
- Morris, R.F. Note on occurrence of the large yellow underwing moth, *Noctua pronuba* (Linnaeus) (Lepidoptera: Noctuidae), in Newfoundland, 403.
- Mukerji, M.K. Parasitism by *Scelio calopteni* Riley (Hymenoptera: Scelionidae) in eggs of the two dominant melanopline species (Orthoptera: Acrididae) in Saskatchewan, 147.
- See Ewen and Mukerji, 537.
- Murvosh, C.M. See Allen and Murvosh, 1095.
- Nealis, V. The number of instars in jack pine budworm, *Choristoneura pinus pinus* Free. (Lepidoptera: Tortricidae), and the effect of parasitism on head capsule width and development time, 773.
- Nealis, V., and J. Régnière. The influence of parasitism by *Apanteles fumiferanae* Vier. (Hymenoptera: Braconidae) on spring dispersal and changes in the distribution of larvae of the spruce budworm (Lepidoptera: Tortricidae), 141.
- Neil, K.A., and H.B. Specht. Sixth-instar larvae of *Noctua pronuba* (L.) (Lepidoptera: Noctuidae), 209.
- Nelson, N. See Turgeon *et al.*, 361.
- Ng, Y.S. See Lashomb *et al.*, 75.
- Niemeyer, H. See McLean *et al.*, 273.
- Nystrom, C.W. See van Frankenhuyzen and Nystrom, 941.
- Obata, S., and T. Hidaka. Ejection and ingestion of the spermatophore by the female ladybird beetle, *Harmonia axyridis* Pallas (Coleoptera: Coccinellidae), 603.
- Onofriechuk, E.E. See Germida *et al.*, 355.
- Oosterbroek, P., and I.R.M. Tangelder. A hybrid population of *Nephrotoma* (Diptera: Tipulidae) from a central Texas refugium, 45.
- A new subspecies, *Nephrotoma lundbecki alexanderi*, described from south of the maximum extent of the Laurentide Ice and new distribution records for other nearctic *Nephrotoma* (Diptera: Tipulidae), 49.
- Osborn, W.P.L., and J.H. Borden. Toxicity of smoke from sporophores of the Indian paint fungus and wood from western hemlock to yellowfever mosquitoes, 1109.

- Osgood, E.A. See Akar and Osgood, 395.
- Otvos, I.S., J.C. Cunningham, and R.I. Alfaro. Aerial application of nuclear polyhedrosis virus against Douglas-fir tussock moth, *Orygia pseudotsugata* (McDunnough) (Lepidoptera: Lymantriidae): II. Impact 1 and 2 years after application, 707.
- Otvos, I.S., J.C. Cunningham, and L.M. Friskie. Aerial application of nuclear polyhedrosis virus against Douglas-fir tussock moth, *Orygia pseudotsugata* (McDunnough) (Lepidoptera: Lymantriidae): I. Impact in the year of application, 697.
- Parry, R.H. See Boiteau *et al.*, 459.
- Peck, S.B., and W.A. Shear. A new blind cavernicolous *Lygromma* (Araneae, Gnaphosidae) from the Galápagos Islands, 105.
- A new eyeless, stridulating *Theridion* spider from caves in the Galápagos Islands (Araneae, Theridiidae), 881.
- Pelletier, Y., and Z. Smilowitz. Specialized tarsal hairs on adult male Colorado potato beetles, *Leptinotarsa decemlineata* (Say), hamper its locomotion on smooth surfaces, 1141.
- Pfadt, R.E. Two new species of *Conophyma* Zubovskii from Afghanistan (Orthoptera: Acrididae), 159.
- Phillips, T.W., S.A. Teale, and G.N. Lanier. Biosystematics of *Pissodes* Germar (Coleoptera: Curculionidae): seasonality, morphology, and synonymy of *P. approximatus* Hopkins and *P. nemorensis* Germar, 465.
- Poirier, D.G., and G.A. Surgeoner. Laboratory flow-through bioassays of four forestry insecticides against stream invertebrates, 755.
- Pree, D.J. See Trimble and Pree, 153.
- Pree, D.J., and H.W. Wagner. Occurrence of cyhexatin and dicofol resistance in the European red mite, *Panonychus ulmi* (Koch) (Acari: Tetranychidae), in southern Ontario, 287.
- Provencher, L. See Coëerre *et al.*, 195.
- Quednau, F.E., and G. Remaudière. Révision des espèces du sous-genre *Castaneomyzocallis* subg. nov. du genre *Myzocallis* trouvées sur les châtaigniers (*Castanea*) en Amérique du Nord (Homoptera: Aphididae), 339.
- Rabasse, J.M. See Lapchin *et al.*, 815.
- Raske, A.G. See West *et al.*, 449.
- See Langor and Raske, 975.
- See Langor and Raske, 985.
- Redner, J.H. See Dondale and Redner, 1.
- Régnière, J. See Nealis and Régnière, 141.
- See Turgeon and Régnière, 239.
- Temperature-dependent development of eggs and larvae of *Choristoneura fumiferana* (Clem.) (Lepidoptera: Tortricidae) and simulation of its seasonal history, 717.
- Rehnberg, B.G. Selection of spider prey by *Trypoxylon politum* (Say) (Hymenoptera: Sphecidae), 189.
- Remaudière, G. See Quednau and Remaudière, 339.
- Retnakaran, A. See West *et al.*, 449.
- Richards, K.W. See Whitfield and Richards, 491.
- See Whitfield *et al.*, 859.
- Ritchie, A.D., and J.D. Shorthouse. Revision of the genus *Synophromorpha* Ashmead (Hymenoptera: Cynipidae), 215.
- Roberts, D.W. See Hajek *et al.*, 959.
- Robinson, A.G. Annotated list of *Macrosiphoniella* (Homoptera: Aphididae) of America north of Mexico, with a key and description of a new species, 913.
- Rutherford, T.A., D. Trotter, and J.M. Webster. The potential of Heterorhabditid nematodes as control agents of root weevils, 67.
- Sánchez, N.E. See Kemp and Sánchez, 653.
- Sanders, C.J., and E.A. Meighen. Controlled-released sex pheromone lures for monitoring spruce budworm populations, 305.
- Schaaflje, G.B. See Byers *et al.*, 557.
- Schmidtman, E.T. A trap/sampling unit for ear-feeding black flies, 747.
- Schulthess, F. See Gutierrez *et al.*, 109.
- Scott-Dupree, C.D., and M.L. Winston. Wild bee pollinator diversity and abundance in orchard and uncultivated habitats in the Okanagan Valley, British Columbia, 735.
- Scudder, G.G.E., and D.B. Thomas, Jr. The green stink bug genus *Chlorochroa* Stål (Hemiptera: Pentatomidae) in Canada, 83.
- Sharkey, M., A. Arthur, G. Bisdee, C. Yoshimoto, and J. Barron. The parasitic Hymenoptera associated with sunflower (*Helianthus* spp.) in mid-western Canada, 611.
- Shear, W.A. See Peck and Shear, 105.
- See Peck and Shear, 881.

- Shemanchuk, J.A.** See Anderson and Shemanchuk, 21.
See Anderson and Shemanchuk, 29.
See Teskey *et al.*, 1117.
- Shipp, J.L.** Diapause induction of eggs of *Simulium arcticum* Malloch (IIS-10.11) (Diptera: Simuliidae), 497.
- Shore, T.L., J.A. McLean, and J.C. Zanuncio.** Reproduction and survival of the ambrosia beetle *Trypodendron lineatum* (Oliv.) (Coleoptera: Scolytidae) in Douglas-fir and western hemlock logs, 131.
- Shorthouse, J.D.** See Ritchie and Shorthouse, 215.
- Smilowitz, Z.** See Pelletier and Smilowitz, 1141.
- Smith, L.B.** See Barker and Smith, 1125.
- Smith, L.B., and P.S. Barker.** Distribution of insects found in granary residues in the Canadian prairies, 873.
- Smith, S.M., M. Hubbes, and J.R. Carrow.** Ground releases of *Trichogramma minutum* Riley (Hymenoptera: Trichogrammatidae) against the spruce budworm (Lepidoptera: Tortricidae), 251.
- Soper, R.S.** See Hajek *et al.*, 959.
- Specht, H.B.** See Neil and Specht, 209.
- Spence, J.R.** See Lee and Spence, 371.
- Stinner, R.E.** See Williams and Stinner, 329.
- Storch, R.H.** See Hajek *et al.*, 959.
- Strongman, D.B.** A method for rearing *Dendroctonus ponderosae* Hopk. (Coleoptera: Scolytidae) from eggs to pupae on host tissue with or without a fungal complement, 207.
- Struble, D.L.** See Byers and Struble, 541.
See Byers *et al.*, 557.
- Struble, D.L., G.L. Ayre, and J.R. Byers.** Sex attractant blends for strawberry cutworm, *Amphipoea inter-oceanica* (Smith), and a closely related species, *Amphipoea americana* (Speyer) (Lepidoptera: Noctuidae), 301.
- Struble, D.L., J.R. Byers, D.G.R. McLeod, and G.L. Ayre.** Sex pheromone components of an Alberta population of European corn borer, *Ostrinia nubilalis* (Hbn.) (Lepidoptera: Pyralidae), 291.
- Surgeoner, G.A.** See Poirier and Surgeoner, 755.
- Sweeney, J.D., and J.A. McLean.** Effect of sublethal infection levels of *Nosema* sp. on the pheromone-mediated behavior of the western spruce budworm, *Choristoneura occidentalis* Freeman (Lepidoptera: Tortricidae), 587.
- Syed, A., and K. Graham.** Response of the mountain pine beetle, *Dendroctonus ponderosae* Hopkins, to ethanol in a laboratory olfactometer, 489.
- Tangelder, I.R.M.** See Oosterbroek and Tangelder, 45.
See Oosterbroek and Tangelder, 49.
- Teale, S.A.** See Phillips *et al.*, 465.
- Teskey, H.J.** See Krzeminski and Teskey, 887.
- Teskey, H.J., J.A. Shemanchuk, and J. Weintraub.** *Hybomitra agora*, a new species of Tabanidae (Diptera) from western North America, 1117.
- Thibodeaux, R.L., W.L. Hutchison, and D.B. Hogg.** Species composition of pea aphid (Homoptera: Aphididae) primary and secondary parasitoids in Wisconsin, 1055.
- Thomas, D.B. Jr.** See Scudder and Thomas, 83.
- Thompson, D.C.** See Capinera and Thompson, 567.
- Tourneur, J.-C.** See Coderre *et al.*, 195.
- Trimble, R.M., and D.J. Free.** Relative toxicity of six insecticides to male and female *Pholetesor ornigis* (Weed) (Hymenoptera: Braconidae), a parasite of the spotted tentiform leafminer, *Phyllonorycter blancardella* (Fabr.) (Lepidoptera: Gracillariidae), 153.
- Trotter, D.** See Rutherford *et al.*, 67.
- Turgeon, J.J., N. Nelson, and E.G. Kettela.** Reproductive biology of the spruce budmoth, *Zeiraphera canadensis* Mut. & Free. (Lepidoptera: Tortricidae: Olethreutinae), in New Brunswick, 361.
- Turgeon, J.J., and J. Régnière.** Development of sampling techniques for the spruce budmoth, *Zeiraphera canadensis* Mut. & Free. (Lepidoptera: Tortricidae), 239.
- Turnock, W.J.** Predicting larval abundance of the bertha armyworm, *Mamestra configurata* Wlk., in Manitoba from catches of male moths in sex attractant traps, 167.
- Turnock, W.J., R.J. Lamb, and R.J. Bilodeau.** Abundance, winter survival, and spring emergence of flea beetles (Coleoptera: Chrysomelidae) in a Manitoba grove, 419.
- van Frankenhuyzen, K.** Effect of wet foliage on efficacy of *Bacillus thuringiensis* spray against the spruce budworm, *Choristoneura fumiferana* Clem. (Lepidoptera: Tortricidae), 955.
- van Frankenhuyzen, K., and C.W. Nystrom.** Effect of temperature on mortality and recovery of spruce budworm (Lepidoptera: Tortricidae) exposed to *Bacillus thuringiensis* Berliner, 941.
- Vickery, V.R.** The orthopteroid insects in northern North America: updating and rationalizing recent work, 1043.
- Villacorta, A.M.** See Gutierrez *et al.*, 109.
- Wagner, H.W.** See Pree and Wagner, 287.
- Wasbauer, M.S.** See Koponen and Wasbauer, 103.

- Weaver, C.A.A. See Fairchild *et al.*, 647.
- Webb, D.W. See Maier and Webb, 55.
- Webster, J.M. See Rutherford *et al.*, 67.
- Weintraub, J. See Teskey *et al.*, 1117.
- Weiss, M.J., and R.J. Beeshear. Seasonal distribution of the flower thrips (*Frankliniella tritici* (Fitch)) and onion thrips (*Thrips tabaci* Lindeman) (Thysanoptera: Thripidae) on safflower in eastern Montana, 399.
- West, R.J., A.G. Raske, A. Retnakaran, and K.P. Lim. Efficacy of various *Bacillus thuringiensis* Berliner var. *kurstaki* formulations and dosages in the field against the hemlock looper, *Lambdina fiscellaria fiscellaria* (Guen.) (Lepidoptera: Geometridae), in Newfoundland, 449.
- Whitc, N.D.G. See Loschiavo and White, 595.
- Whitfield, G.H., and K.W. Richards. Postdiapause development and adult emergence of *Pteromalus venustus* Walker (Hymenoptera: Pteromalidae) during alfalfa leafcutter bee incubation, 491.
- Whitfield, G.H., K.W. Richards, and T.M. Kveder. Number of instars of larvae of the alfalfa leafcutter bee, *Megachile rotundata* (F.) (Hymenoptera: Megachilidae), 859.
- Whiting, E.R., and D.M. Lehmkuhl. *Raptoheptagenia cruentata*, gen. nov. (Ephemeroptera: Heptageniidae), new association of the larva previously thought to be *Anepeorus* with the adult of *Heptagenia cruentata* Walsh, 405.
- Acanthomola pubescens*, a new genus and species of Heptageniidae (Ephemeroptera) from western Canada, 409.
- Wiggins, G.B., and N.W. Erman. Additions to the systematics and biology of the caddisfly family Uenoidae (Trichoptera), 867.
- Williams, D.W., and R.E. Stinner. Respiration and drowning in overwintering pupae of *Heliothis zea* (Boddie), 329.
- Wilson, L.T. See Gutierrez *et al.*, 109.
- Winston, M.L. See Scott-Dupree and Winston, 735.
- Wong, H.R. See Wu and Wong, 397.
- Wright, B. The European yellow underwing, *Noctua pronuba* L. (Lepidoptera: Noctuidae), in the Atlantic Provinces (Canada) and the state of Maine (USA), 993.
- Wu, J., and H.R. Wong. Colonization of lodgepole pine stumps by ants (Hymenoptera: Formicidae), 397.
- Wu, K.W. See Lindquist and Wu, 779.
- Wylie, H.G., and H.E. Bindee. Primary and secondary parasites of an alfalfa-infesting aphid, *Therioaphis* sp., in Manitoba, 857.
- Wynne, A.D. See Javer *et al.*, 605.
- Yoshimoto, C. See Sharkey *et al.*, 611.
- Young, T.-L., R. Gordon, and M. Cornect. Effects of several insect growth regulators on egg hatch and subsequent development in the cabbage maggot *Delia radicum* (L.) (Diptera: Anthomyiidae), 481.
- Yu, D.S.K. See Jaques *et al.*, 1063.
- Zanuncio, J.C. See Shore *et al.*, 131.

